Prolonged Standing and Work Fatigue on Sales Promotion Girl (SPG) Workers at the Kubu Raya Regency Shopping Center

by Sunarsieh Sunarsieh

Submission date: 29-Mar-2023 09:17AM (UTC+0700)

Submission ID: 2049581777

File name: 22677-44131-1-RV an Sunarsieh.docx (122.82K)

Word count: 3902 Character count: 21444 ISSN: 2252-8806, DOI: 10.11591/ijphs.vxix.id

101

Prolonged Standing and Work Fatigue on Sales Promotion Girl (SPG) Workers at the Kubu Raya Regency Shopping Center

Sunarsieh¹, Moh Adib ¹

¹Department of Environmental Health, Poltekkes Kemenkes Pontianak, West Borneo, Indonesia

Article Info

Article history:

Received Revised Accepted

Keywords:

Length of working Prolonged standing Work fatigue Working position standing SPG Workers

ABSTRACT

Standing positions are used for almost all jobs, including Sales Promotion Girl workers. Standings in working positions for long periods can cause work fatigue. This study aims to analyze the relationship between the length of work in a standing position and the work fatigue of SPG workers. This research is a cross-sectional study. Fatigue data collection was carried out by interviews using the Industrial Fatigue Research Committee (IFRC) questionnaire. Data on the length of work in a standing position were obtained by interview and observation. Based on the results of the study, as many as 66.7% of workers at SPG in the shopping center of Kubu Raya Regency in one day work in a standing position for 8 hours, and 33.3% work in a standing position for >8 hours. The average level of work fatigue at the beginning of work is 49.38, and after work is 52.95, with the results of statistical analysis showing a p-value of 0.000, meaning that there is a significant relationship between long-standing position and work fatigue. There is a need for supervision and monitoring related to the work safety system regularly carried outaby the manager of the shopping center, which is also supervised by the Ministry of Health and the Ministry of Manpower.

This is an open access article under the CC BY-SA license.



Corresponding Author:

Sunarsieh

Department of Environmental Health, Poltekkes Kemenkes Pontianak, West Borneo, Indonesia Jl.28 Oktober, Siantan Hulu, Pontianak, West Borneo, Indonesia

Email: asiehbima@gmail.com

1. INTRODUCTION

Occupational health and safety efforts aim to protect workers so that they live healthy and free from health problems, occupational diseases, and adverse effects caused by work [1]. Ergonomic factors such as an uncomfortable working position and working in a prolonged standing position can result in occupational diseases [2]. Occupational diseases are diseases caused by exposure to risk factors arising from work activities, which can be caused by work environment factors and other risk factors such as ergonomic factors, such as prolonged standing positions [3]. The standing position is used in almost all jobs, including Sales Promotion Girl (SPG) in shopping centers. Sales Promotion Girl performs work activities in a prolonged standing position for most of her working hours. SPG workers generally spend 50% of their total working hours in a prolonged standing position [4]. SPG workers also have to work long hours by lifting activities, carrying goods, serving customers, offering products, doing repetitive movements, and standing for long periods [5]. Prolonged standing during work has a high risk of causing chronic or acute health problems. Standing for too long reduces the muscles' blood supply, resulting in fatigue [6], [7]. According to (CCOHS, 2014), fatigue can impact work accidents. As many as 13% of work accidents are caused by work fatigue, and as many as 16% of workers experience work accidents at least once due to fatigue [8]. Of the number of accidents that occur in the

Journal homepage: http://ijphs.iaescore.com

workplace, 60% are caused by work fatigue. [9], [10] Accumulated worldwide A total of 6000 fatal accidents resulting in many victims [2].

The observations made by researchers at shopping centers in Kubu Raya Regency showed SPG workers work in a standing position during their working hours and the result of interviews with SPG workers work 8 hours a day. Workers are exposed to prolonged standing because they spend more than 50% of their total working hours in a standing position. Based on these problems, the researchers conducted a study on the length of standing work positions and work fatigue on SPG workers at the Kubu Raya Regency Shopping Center.

The purpose of this study is to examine the standing working position, know the long-standing working position, assess fatigue at the start of work and after work, analysis of the relationship between the standing position and work fatigue of SPG workers. This research is necessary because physical and ergonomic factors such as standing work positions need attention and are a common problem in the workplace [11]. The prevalence and severity of health problems are significantly higher in employees who work in a standing position for a long time. Hence, SPG workers need to be research subjects so that later interventions can be carried out [12].

2. RESEARCH METHOD

This type of research is cross-sectional and aims to determine the relationship between the length of work in a standing position and work fatigue in SPG workers. A total of 60 SPG workers were involved in this study as research respondents. This research has obtained ethical approval from the Health Research Ethics Committee of the Health Polytechnic of the Pontianak Ministry of Health. No. 50/KEPK-PK.PKP/III/2022. Research respondents willing to participate in the study have signed an informed consent form. The research was conducted April-September 2022. SPG workers involved in this study have met the inclusion and exclusion criteria. Data collection on the length of work standing was collected using interviews and observations using questionnaires and checklists. Work fatigue data collection uses the IFRC (International Fatigue Research Committee of the Japanese Association of Industrial Health) questionnaire. The IFRC questionnaire contains 30 general fatigue questions used to measure fatigue. The first ten questions on the IFRC questionnaire indicate a weakening of activity; the second 10 questions indicate a decline in work motivation. And the last ten questions indicate fatigue in several body parts or physical fatigue in the respondent. After conducting interviews and collecting respondent data, the next step is to calculate the total score from the answers to the 30 questions organized. The total number becomes the total individual score. This questionnaire was then developed, and the answers to the questionnaire were scored according to four Likert scales. Based on the results of the fatigue assessment, of the 30 questions, the highest score was 120. The answers to the IFRC questionnaire were divided into four categories, namely Very Often (SS) with a score of 4, Often (S) with a value of 3, Sometimes (K) with a score of 2, and Never (T) with a value of 1. In determining the fatigue classification, the answers to each question are added and then adjusted to a certain classification. The classifications given include: Score >55 = High Fatigue and Score 55 = Low Fatigue [13], [14].

The procedure in this study starts from the preparation stage for the management of research ethics approval, secondary data collection, surveys, and field observations to identify problems, research permits, and meetings between the research team and data collection officers for the division of tasks and research perceptions. The next stage of implementation starts with surveys, field observations, and inventory of work activities carried out by SPG workers, meetings with companies to explain the procedures for research activities, determining research respondents, explaining research procedures and actions to respondents, collecting respondent data and collecting work fatigue data before and after work.

The research data were then processed and analyzed by descriptive analysis to determine the distribution characteristics of each dependent and independent variable. The activities of this descriptive statistical analysis include presenting data using tables and graphs with simple calculations such as percentages, averages, and ratios. Furthermore, inferential analysis is carried out using measures with estimation theory. They tested the research hypothesis using the chi-square statistical test using a 95% confidence level so that the relationship between research variables could be known. The test was carried out with a significance level (α = 0.05). If p < 0.05, then the test results were significant.

3. RESULTS AND DISCUSSIONS

Table 1. Demographic Distribution of SPG Workers

Variable	Category		
		n	%
Marital status	Single	48	80
	Marry	12	20
Age	< 25 years	46	76.7
	≥25 years	14	23.3
Education	Basic education	1	1.7
	Middle education	57	95
	Higher education	2	3.3
Working period	< 1 years	16	26.7
	1-5 years	35	58.3
	> 5 years	9	15
Working length	<= 8 hours/day	40	66.7
	> 8 hours/day	20	33.3

Table 1 shows the demographic distribution of SPG Shopping Center workers in Kuburaya Regency. Most of the respondents were unmarried, as many as 48 people (80%). A total of 46 people (76.7%) of respondents were in the age category <25 years. A total of 35 people (58.3%) of respondents have worked for 1-5 years. A total of 40 people (66.7%) of respondents work 8 hours a day.

Table 2. The Results of The Work Fatigue analysis of SPG Workers

Work Fatigue	Category	n	Mean±sd	CI = 95%		p-value
				min	max	
Fatigue before and after	Before work	60	49,38 <u>+</u> 8,876	35	67	0.000
work	After work	60	52.95+10.834	35	87	

Table 2 shows the results of the work fatigue analysis, namely the comparison between the total work fatigue scores before and after working for SPG workers at the Kuburaya Regency Shopping Center with a pvalue of 0.000 which means there is a significant difference in work fatigue before and after working for SPG workers at the Regency Shopping Center. Burial. Based on the results of the study, the relationship between the length of work in a standing position and work fatigue on SPG workers at the Kuburaya Regency Shopping Center with a p-value of 0.000. Fatigue experienced by a worker is characterized by decreased morale due to work that is done too monotonously, work that is charged too much, demands for fast work completion times, work postures that are not ergonomic, nutritional status is not normal, psychological conditions of workers, age factors, habits breakfast and how long a person's experience is doing his job [15]. Fatigue at work is caused by the interaction of many factors, including shift patterns, physical workload, monotonous and unstimulating work, poor sleep habits, and other lifestyle factors. Fatigue can lead to unwanted accidents resulting in reduced work productivity and a significant decline in decision-making skills and memory acuity. [16] [17]. Fatigue before work is a sign of someone experiencing chronic fatigue. Fatigue before work is caused by activities carried out by someone that has been going on for a long time and appears throughout the day before doing work activities, causing difficulty sleeping, pain in the head, and digestive problems [18]. Work fatigue before work is caused by work environment, work type, individuals, and work positions. [19], [20]. Prolonged standing positions increase fatigue throughout the body, so management of adequate rest time can help reduce work fatigue [20]. 33.3% of SPG workers at the Kuburaya Regency Shopping Center work more than 8 hours. Standard workers generally work 8 hours a day in a prolonged standing position. However, if workers work more than the maximum working hours with a prolonged standing position, it can cause workers to experience muscle fatigue [21]. Standing for more than 5 hours can cause muscle fatigue that lasts more than 30 minutes after work hours [22]. The addition of working hours in a non-ergonomic position can also affect the ability and trigger fatigue in workers. If done continuously can cause chronic fatigue [23].

Table 3. The Relationship of Length of Work in Standing Work Positions with Work Fatigue in SPG Workers

Variable	Category	n	CI = 95 %		r count	
			Mean	SD	(p-value)	
Long working relationship with a	Length of working	60	9,00	1,584	0,322 (0.010)	
standing position with fatigue after work	Fatigue after work	60	52,95	10,834		

Table 3 shows the results of the analysis of the relationship between the length of work in a standing position and work fatigue for SPG workers at the Kubu Raya Regency Shopping Center, the p-value of 0.010, which means there is a relationship between the length of work in a standing position and work fatigue after work. SPG workers generally work in a standing position for hours and even exceed the standing time limit, so they have a great potential to experience work fatigue. According to Garcia et al., the standing duration can cause muscle fatigue in the back and lower extremities [22]. Work that requires long-standing causes discomfort because it requires workers to try to balance their body position and causes a static workload in the back muscles, legs, and physical fatigue [20]. SPG (Sales Promotion Girl) must stand for a long time, for 7-8 hours per day. Ergonomic working positions can reduce the risk of injury to workers. However, working postures carried out for a long time without stretching can cause leg circulation problems in blood circulation, swelling in congenital diseases, and cause fatigue located in the leg muscles [24]. Standing for extended periods can cause fatigue in workers, triggering headaches and increasing susceptibility to disease. [25]. The intervention has an optional chair or footrest to increase the variety of body positions available to standing workers. [26]. In addition to footrests, other interventions are ergonomic training programs for workers regularly and job rotations [27]. Standing interspersed with sitting for a while is an excellent solution to minimize work fatigue associated with the working position standing at work when part of the muscles works lightly to stabilize the body that is experiencing fatigue [28]. However, sitting for a long time can also affect health, so changing work positions and sitting using a chair/bench is necessary [7], [29]. Another intervention to reduce work fatigue is continuous muscle contraction to maintain a prolonged posture can increase whole-body fatigue. Rest and stretching can reduce overall body fatigue. [20]. Work posture is a determining point in analyzing the effectiveness of a job. If the work posture carried out by the worker is excellent and ergonomic, the results obtained by the worker will certainly be good. However, if the work posture is not good, the worker will easily experience fatigue [30]. Occupational health and safety monitoring work to protect workers from hazards, improve occupational health and safety in the work environment, and prevent accidents and occupational diseases. The purpose of monitoring worker health and safety is to ensure that occupational health and safety programs are implemented effectively [31]. Monitoring of occupational health and safety must also be carried out periodically to find out that the implementation of the occupational health and safety system in the company is running effectively. Monitoring a company's occupational health and safety system can reduce the risk of health problems and accidents, increase productivity and reduce unexpected company expenses [32]. Occupational health and safety monitoring activities are carried out by auditing the occupational health and safety management system. The occupational health and safety management system audit aims to protect workers from occupational diseases and the risk of work accidents, increase productivity, minimize risks, and improve the company's image [33]. OHS management system audits are conducted periodically to determine the effectiveness of implementing the OHS management system. Audits are carried out independently and systematically by personnel/institutions with the appropriate competencies. The frequency of audits is determined based on a review of the results of previous audits and evidence of identified hazard sources in the workplace environment. The management should use the results in the management review process. Audits are carried out two times a year and involve all parts of the company in each section/field. [34].

4. CONCLUSION

This study revealed that as many as 66.7% of workers at SPG in the shopping center of Kubu Raya Regency one day work in a position for 8 hours, and 33.3% work in a standing position for>8 hours. The average level of work fatigue at the beginning of work is 49.38, and after work is 52.95, with the results of statistical analysis showing a p-value of 0.000. Based on the statistical analysis results, there is a relationship between the length of work in a standing position and fatigue after working for SPG workers at the Kubu Raya Shopping Center, with a p-value of 0.010. From the results of this study, it is necessary to have supervision and monitoring

related to the work safety system regularly carried out by the management of the shopping center, which is also supervised by the Ministry of Health and the Ministry of Manpower.

ACKNOWLEDGEMENTS

We would like to thank Poltekkes Kemenkes Pontianak. And we also thank the mall management, data collection officers, and SPG workers who were research participants for their cooperation during this research activity.

REFERENCES

- [1] WHO, "Protecting workers health," World Health Organization, 2017. .
- [2] International Labor Organization, Meningkatkan Keselamatan dan Kesehatan Pekerja Muda. Jakarta: International Labor Organization Switzerland, 2018.
- [3] WHO, "Occupational Health," World Health Organization, 2017. https://www.who.int/health-topics/occupational-health (accessed Aug. 01, 2022).
- [4] E. M. Capodaglio, "Occupational risk and prolonged standing work in apparel sales assistants," Int. J. Ind. Ergon., vol. 60, pp. 53–59, 2017, doi: 10.1016/j.ergon.2016.11.010.
- [5] R. S. Purani, N. J. Vyas, and M. S. Sheth, "Prevalence of low back pain in salespersons and its association with ergonomic risk factors in Ahmedabad, Gujarat: A cross-sectional survey," Med. J. Dr. D.Y. Patil Vidyapeeth, vol. 9, no. 3, pp. 331–335, 2016, doi: 10.4103/0975-2870.182503.
- [6] D. Antle, N. Vezina, K. Messing, and J. Cote, "Development of discomfort and vascular and muscular changes during a prolonged standing task," *Occup. Ergon.*, vol. 11, no. 1, pp. 21–33, 2013, doi: 10.3233/OER-130205.
- [7] T. R. Waters and R. B. Dick, "Evidence of health risks associated with prolonged standing at work and intervention effectiveness," *Rehabil. Nurs.*, vol. 40, no. 3, pp. 148–165, 2015, doi: 10.1002/rnj.166.
- [8] CCOHS, "Work related musculoskeletal disorders (WMSDs)," Canadian Centre for Occupational Health and Safety, 2014. https://www.ccohs.ca/oshanswers/diseases/rmirsi.html#:~:text=What are work-related musculoskeletal,tension neck syndrome are examples.
- [9] K. Uehli et al., "Sleep problems and work injuries: A systematic review and meta-analysis," Sleep Med. Rev., vol. 18, no. 1, pp. 61–73, 2014, doi: 10.1016/j.smrv.2013.01.004.
- [10] Safe Work Australia, Guide For Managing The Risk Of Fatigue At Work. 2013.
- [11] L. R. Prado-León and E. Herrera-Lugo, "Manual Lifting Standards: Ergonomic Assessment and Proposals for Redesign for Industrial Applications," in New Perspectives on Applied Industrial Tools and Techniques. Management and Industrial Engineering, Springer, 2018.
- [12] Z. Karimi et al., "Determining the interactions between postural variability structure and discomfort development using nonlinear analysis techniques during prolonged standing work," Appl. Ergon., vol. 96, no. June, p. 103489, 2021, doi: 10.1016/j.apergo.2021.103489.
- [13] I. M. Ramdan, "Measuring Work Fatigue on Nurses: A Comparison between Indonesian Version of Fatigue Assessment Scale (FAS) and Japanese Industrial Fatigue Ressearch Commite (JIFRC) Fatigue Questionnaire," J. Keperawatan Padjadjaran, vol. 7, no. 2, pp. 143–153, 2019, doi: 10.24198/jkp.v7i2.1092.
- [14] T. Ihsan, T. Edwin, Y. Azwir, and V. Derosya, "Fatigue Analysis to Evaluate Workloads in Production Area at Crumb Rubber Factories of Padang city, West Sumatra Indonesia," *Indian J. Occup. Environ. Med.*, vol. 24, no. 3, pp. 148–152, 2020, doi: 10.4103/ijoem.IJOEM_252_19.
- [15] D. C. Dewi, J. Hutabarat, and P. Vitasari, "Pengukuran Kelelahan Shift Kerja pada Pekerja Pabrik Kelapa Sawit PT. BGR Jambi," J. Teknol. dan Manaj. Ind., vol. 2, no. 1, pp. 21–27, 2016.
- [16] RoSPA, "Fatigue and Work: RoSPA Position Statement," Birmingham, 2019.
- [17] S. Bendak and H. S. J. Rashid, "Fatigue in aviation: A systematic review of the literature," Int. J. Ind. Ergon., vol. 76, p. 102928, 2020, doi: 10.1016/j.ergon.2020.102928.
- [18] Centers for Disease Control and Prevention, "Myalgic Encephalomyelitis/ Chronic Fatigue Syndrome," U.S.Department of Health & Human Services, 2022.
- [19] J. A. Caldwell, J. L. Caldwell, L. A. Thompson, and H. R. Lieberman, "Fatigue and its management in the workplace," *Neurosci. Biobehav. Rev.*, vol. 96, no. November 2018, pp. 272–289, 2019, doi: 10.1016/j.neubiorev.2018.10.024.
- [20] H. Jo, O. Bin Lim, Y. S. Ahn, S. J. Chang, and S. B. Koh, "Negative impacts of prolonged standing at work on musculoskeletal symptoms and physical fatigue: The fifth korean working conditions survey," *Yonsei Med. J.*, vol. 62, no. 6, pp. 510–519, 2021, doi: 10.3349/ymj.2021.62.6.510.
- [21] N. R. Hauteas, L. P. Ruliati, and S. Doke, "Analysis of Factors Affecting Work Fatigue on Wig Workers at CV. Dona Mandiri Lasiana Branch Kupang City," *J. Community Health*, vol. 1, no. 4, 2019, doi: https://doi.org/10.35508/ljch.v1i4.2178.

106 □ ISSN: 2252-8806

[22] M. G. Garcia, T. Läubli, and B. J. Martin, "Long-term muscle fatigue after standing work," Hum. Factors, vol. 57, no. 7, pp. 1162–1173, 2015, doi: 10.1177/0018720815590293.

- [23] L. Sulistioningsih, "Factors Associated with Work Fatigue on Workers in the Food Production Section," *Medica Majapahit*, vol. 5, no. 1, 2013, doi: 10.5220/0009785103920396.
- [24] S. M. Anggrianti, B. Kurniawan, and B. Widjasena, "Hubungan Antara Postur Kerja Berdiri dengan Keluhan Nyeri Kaki pada Pekerja Aktivitas Mekani Section Welding di PT. X," J. Kesehat. Masy., vol. 5, no. 5, pp. 369–377, 2017.
- [25] CCOHS, "Anti Fatigue Mats," Canadian Centre for Occupational Health and Safety, 2015. .
- [26] Canadian Centre for Occupational Health and Safety, "OSH Answers Fact Sheet: Fatigue," Government of Canada, 2017. https://www.ccohs.ca/oshanswers/psychosocial/fatigue.html.
- [27] R. Heidarimoghadam, I. Mohammadfam, M. Babamiri, A. R. Solatanian, H. Khotanlou, and M. S. Sohrabi, "What do the different ergonomic interventions accomplish in the workplace? A systematic review," *Int. J. Occup. Saf. Ergon.*, vol. 28, no. 1, pp. 600–624, 2022, doi: 0.1080/10803548.2020.1811521.
- [28] D. M. Antle, L. Cormier, M. Findlay, L. L. Miller, and J. N. Côté, "Lower limb blood flow and mean arterial pressure during standing and seated work: Implications for workplace posture recommendations," *Prev. Med. Reports*, vol. 10, no. October 2017, pp. 117–122, 2018, doi: 10.1016/j.pmedr.2018.02.016.
- [29] T. J. Saunders, H. F. Atkinson, J. Burr, B. MacEwen, C. M. Skeaff, and M. C. Peddie, "The Acute Metabolic and Vascular Impact of Interrupting Prolonged Sitting: A Systematic Review and Meta-Analysis," Sport. Med., vol. 48, no. 10, pp. 2347–2366, 2018, doi: 10.1007/s40279-018-0963-8.
- [30] I. Yudisianto, A. R. Tualeka, and N. Widajati, "Correlation between Individual Characteristics and Work Position with Work Fatigue on Workers," *Indones. J. Occup. Saf. Heal.*, vol. 10, no. 3, p. 350, 2021, doi: 10.20473/ijosh.v10i3.2021.350-360.
- [31] International Labour Organization, Quick Guide on Sources and Uses of Statistics on Occupational Safety and Health, 1st ed. 2020.
- [32] A. Zikriyoev, "The Business Benefits of Good Occupational Health and Safety in Construction Industry," Asian J. Multidimens. Res., vol. 8, no. 3, 2019, doi: 10.5958/2278-4853.2019.00129.0.
- [33] M. A. Y. Saputra, M. N. Fadhillah, and D. Setiadi, "Aplikasi Checklist Safety Berbasis Mobile Menggunakan Metode Job Safety Analysis (JSA) Untuk Audit Kesehatan Dan Keselamatan Kerja (K3) Di Pt.Matahari Putra Prima, Tbk," J. Sist. Inf., pp. 257–261, 2021.
- [34] A. A. S. Budi and A. S. Wahyuningsih, "Aspek Keselamatan dan Kesehatan Kerja di Supermall," Higeia J. Public Heal. Res. Dev., vol. 1, no. 3, pp. 625–634, 2019.

BIOGRAPHIES OF AUTHORS



Sunarsieh, a graduate of Hyperkes and Occupational Health and Safety in the Doctoral Program of Universitas Gadjah Mada, Yogyakarta, 2013. She has served as a lecturer in the Environmental Health Department of the Poltekkes Kemenkes Pontianak with the position of associate professor. As well as being a lecturer, she has been given the mandate as Deputy Director in the field of Student Affairs since 2018. And she is currently a member of the Association of Environmental Health Experts (HAKLI) of West Kalimantan Province and Member of the Association of Indonesian Occupational Health Experts (PAKKI) West Kalimantan.



Moh Adib, graduated from Public Health Sciences at the Masters Program at the University of Indonesia, Jakarta, 2000. He currently serves as a lecturer at Poltekkes, Ministry of Health, Pontianak with the position of Assistant Professor. Apart from being a lecturer, he has been given the mandate of Chairman I of The Indonesian Public Health Association/IPHA of West Kalimantan Province.

Prolonged Standing and Work Fatigue on Sales Promotion Girl (SPG) Workers at the Kubu Raya Regency Shopping Center

ORIGINALITY REPORT				
SIMIL	5% 13% INTERNET SOURCE	6% DES PUBLICATIONS	3% STUDENT PAPERS	
PRIMAF	RY SOURCES			
1	kesma.undana.ac.id		2%	
2	ieomsociety.org Internet Source		1 %	
3	conference.juriskes.o	com	1 %	
4	Thomas A. Hethmon "Industrial Hygiene A Hygiene", Wiley, 200	Abroad: Occupati	0/-	
5	ejournal.unsrat.ac.id		1 %	
6	repo.stikesicme-jbg.	ac.id	1 %	
7	Submitted to School Management ITB Student Paper	of Business and	1 %	
8	jurnal.stikescirebon.	ac.id	1 %	

9	www.rospa.com Internet Source	1 %
10	cdu.herdin.ph Internet Source	1 %
11	Isa Halim, Abdul Rahman Omar. "Development of Prolonged Standing Strain Index to Quantify Risk Levels of Standing Jobs", International Journal of Occupational Safety and Ergonomics, 2015 Publication	<1%
12	Submitted to Universitas Airlangga Student Paper	<1%
13	"Occupational and Environmental Safety and Health III", Springer Science and Business Media LLC, 2022 Publication	<1%
14	Gosia Speed, Kate Harris, Tessa Keegel. "The effect of cushioning materials on musculoskeletal discomfort and fatigue during prolonged standing at work: A systematic review", Applied Ergonomics, 2018 Publication	<1%
15	repository.upnvj.ac.id Internet Source	<1%
16	knowledgecommons.popcouncil.org	<1%

17	www.researchgate.net Internet Source	<1%
18	www.semanticscholar.org Internet Source	<1 %
19	jhsw.tums.ac.ir Internet Source	<1%
20	link.springer.com Internet Source	<1%
21	ojs.uho.ac.id Internet Source	<1%
22	plone.ufpb.br Internet Source	<1 %
23	theses.gla.ac.uk Internet Source	<1 %
24	www.ncbi.nlm.nih.gov Internet Source	<1%

Exclude quotes On Exclude bibliography On

Exclude matches

Off

Prolonged Standing and Work Fatigue on Sales Promotion Girl (SPG) Workers at the Kubu Raya Regency Shopping Center

GRADEMARK REPORT	
FINAL GRADE	GENERAL COMMENTS
/0	Instructor
,	
PAGE 1	
PAGE 2	
PAGE 3	
PAGE 4	
PAGE 5	
PAGE 6	
PAGE 7	